

Supplementary Table 4. Diagnostic changes of all participants

Subject number	Initial	After 1 year	Treatment	EBT
1	Adjustment disorder	No diagnosis	Play therapy, psychotherapy	O
2	PTSD	PTSD Depressive disorder NOS	Psychotherapy	O
3	Depressive disorder NOS	No diagnosis	Psychotherapy, TF-CBT	O
4	No diagnosis	No diagnosis	-	
5	PTSD	No diagnosis	-	
6	PTSD	PTSD	TF-CBT	O
7	PTSD Depressive disorder NOS	PTSD Depressive disorder NOS	Medication	O
8	PTSD	No diagnosis	Medication	O
9	Depressive disorder NOS	No diagnosis	TF-CBT	O
10	Depressive disorder NOS	No diagnosis	Psychotherapy TF-CBT Medication	O
11	No diagnosis	No diagnosis	-	
12	Depressive disorder NOS	No diagnosis	-	
13	Enuresis	Depressive disorder NOS		
14	PTSD	PTSD	Medication	O
15	Adjustment disorder	No diagnosis	-	
16	PTSD	No diagnosis	Psychotherapy	O
17	PTSD Depressive disorder NOS	PTSD	Medication	O
18	PTSD Depressive disorder NOS	PTSD	TF-CBT	O
19	PTSD	ADHD	Psychotherapy TF-CBT EMDR	O
20	Depressive disorder NOS	No diagnosis	-	
21	No diagnosis	No diagnosis	Art therapy	
22	No diagnosis	No diagnosis	Art therapy	
23	PTSD	PTSD	Medication	O
24	PTSD	PTSD	Medication	O
25	Anxiety disorder NOS	No diagnosis	-	
26	PTSD	PTSD	Medication	O
27	No diagnosis	No diagnosis	-	
28	PTSD	No diagnosis	Medication	O

Psychotherapy means from supportive psychotherapy to psychodynamic psychotherapy. Medication means that the subject used one or more of the following drugs; selective serotonin reuptake inhibitors, antipsychotics adjuvant, serotonin and norepinephrine reuptake inhibitor, tricyclic antidepressants, and mood stabilizers. EBT, evidence-based treatment; PTSD, posttraumatic stress disorder; NOS, not otherwise specified; ADHD, attention-deficit/hyperactivity disorder; TF-CBT, trauma-focused cognitive behavior therapy; EMDR, eye movement desensitization and reprocessing