

Supplementary Table 1. Characteristics of suicidal ideation among confirmed COVID-19 patients

	Suicidal ideation		t-value λ^2	p
	Positive (weighted N=19)	Negative (weighted N=621)		
Sex			1.96	0.162
Male	7 (35.0)	316 (50.9)		
Female	13 (65.0)	305 (49.1)		
Age (yr)	50.16±16.57	46.53±16.32	0.95	0.352
Marital status			0.07	0.966
Unmarried	6 (31.6)	200 (32.2)		
Married	10 (52.6)	336 (54.1)		
Divorced/separated/widowed	3 (15.8)	85 (13.7)		
Employment status			0.18	0.674
Employed	12 (63.2)	420 (67.7)		
Unemployed	7 (36.8)	200 (32.3)		
Educational status			6.12	0.013
High school graduation or lower	6 (31.6)	372 (59.9)		
College graduation or higher	13 (68.4)	249 (40.1)		
Family income (million won)			10.13	0.018
<1.5	6 (30.0)	55 (8.9)		
1.5–2.99	4 (20.0)	157 (25.3)		
3–4.99	4 (20.0)	182 (29.3)		
≥5	6 (30.0)	227 (36.6)		
COVID-19 vaccination	14 (70.0)	546 (87.9)	5.64	0.018
Rating scales scores				
Health satisfaction of COVID-19 survivors	7.69±2.20	5.11±2.37	4.74	<0.001
Sleep time	6.09±1.77	6.92±1.10	2.06	0.056
PSQI-K	9.43±5.59	4.08±2.62	4.22	<0.001
CFS-K	21.05±8.07	12.09±5.04	4.87	<0.001
Stigma index	21.88±6.19	19.33±5.72	1.93	0.054
Loneliness index	7.99±3.11	5.43±2.62	4.21	<0.001

Values are presented as number (%) or mean±standard deviation unless otherwise indicated. COVID-19, coronavirus disease-2019; PSQI-K, Pittsburgh Sleep Quality Index (total score 21); CFS-K, Chronic Fatigue Scale (total score 33)