

Supplementary Table 2. Raw scores of the EORTC QLQ-CIPN20 scales of groups by PSS from friends, by time point

	Time point	Low-to-moderate PSS		High PSS	
		N	Mean±SD	N	Mean±SD
Total score	T0	12	5.71±8.28	27	5.03±7.29
	C3	10	16.45±16.64	26	6.51±7.45
	C7	9	30.01±13.65	19	15.97±10.51
	T1	10	30.38±18.11	24	17.99±17.21
	T2	4	4.00±3.61	11	13.57±15.94
Sensory scale	T0	12	4.01±6.41	27	4.53±6.52
	C3	10	17.41±16.38	26	4.99±9.01
	C7	9	38.68±20.63	19	20.86±18.44
	T1	10	36.67±18.84	24	20.99±22.72
	T2	4	5.56±8.82	11	15.49±21.90
Motor scale	T0	12	4.46±8.26	27	3.37±7.89
	C3	10	10.77±17.75	26	5.20±6.16
	C7	9	17.13±14.20	19	7.86±9.11
	T1	10	21.55±19.14	24	13.54±16.50
	T2	4	1.19±2.38	11	8.98±10.19
Autonomic scale	T0	12	18.06±26.07	27	13.58±20.69
	C3	10	33.33±27.22	26	18.59±19.05
	C7	9	40.74±20.60	19	25.44±21.78
	T1	10	36.67±33.15	24	21.53±19.95
	T2	4	8.33±9.62	11	22.73±20.10

EORTC-QLQ CIPN20, European Organization for Research and Treatment of Cancer Quality of Life Questionnaire-Chemotherapy-Induced Peripheral Neuropathy 20; PSS, perceived social support; SD, standard deviation