Supplementary Table 2. Stratified analysis of CBT outcomes by baseline cognitive ability (FSIQ), attendance rate, and comorbid depression

Stratification factors		ASRS			K-AARS							
		Part A	Part B	Total	Inattention	Hyperactivity	Impulsivity	APD/CD/ODD	Dysregulation	Disorganization	Subtotal	Impairment
FSIQ												
≥110	β*	-1.600	-4.400	-6.000	-6.800	-0.600	-1.800	-1.600	-2.200	-2.100	-15.100	-2.500
	p-value	0.037	0.008	0.002	0.037	0.329	0.188	0.029	0.181	0.079	0.010	0.024
90–109	β*	-4.222	-7.889	-12.111	-14.000	-2.556	-7.667	-3.333	-11.444	-5.667	-44.667	-4.778
	p-value	0.010	0.006	0.004	<0.001	0.071	<0.001	0.002	<0.001	0.001	<0.001	0.004
<90	β*	-6.000	-11.250	-17.250	-16.000	0.750	-7.750	-2.000	-10.500	-3.250	-38.750	-3.750
	p-value	0.038	0.072	0.057	0.049	0.319	0.023	0.041	0.006	0.152	0.023	0.312
Attendance rate												
100%	β*	-3.846	-7.000	-10.846	-13.615	-1.539	-5.231	-2.000	-7.308	-5.539	-35.231	-3.846
	p-value	0.003	0.008	0.004	0.000	0.132	0.001	0.014	0.001	<0.001	<0.0001	0.011
<100%	β*	-2.800	-6.900	-9.700	-8.100	-0.600	-5.000	-2.800	-7.200	-1.300	-25.000	-3.300
	p-value	0.013	<0.0001	0.000	0.006	0.343	0.015	<0.0001	0.020	0.146	0.004	0.002
Comorbid depression												
Depression	β*	-3.727	-7.818	-11.546	-13.273	-1.091	-6.455	-2.636	-9.364	-4.546	-37.364	-4.455
	p-value	0.006	0.006	0.004	0.000	0.238	0.000	0.005	0.001	0.002	<0.0001	0.006
Non-comorbid	β*	-3.083	-6.167	-9.250	-9.333	-1.167	-3.917	-2.083	-5.333	-2.917	-24.750	-2.833
	p-value	0.009	0.001	0.001	0.005	0.198	0.022	0.002	0.017	0.015	0.002	0.014

*estimated effects adjusted for baseline covariates including age, sex, education, occupation, marital status, comorbidity, medication, and K-WAIS-IV. CBT, cognitive behavioral therapy; FSIQ, full-scale intelligence quotient; ADHD, attention-deficit/hyperactivity disorder; ASRS, Adult ADHD Self-Report Scale; K-AARS, Korean Adult ADHD Rating Scale; APD/CD/ODD, antisocial personality disorder/conduct disorder/oppositional defiant disorder; K-WAIS-IV, Korean Wechsler Adult Intelligence Scale-IV.