Supplementary Table 1. Predictors of clinically significant depressive, anxiety, and PTSD symptoms at 2 months post-incident†

	Depression			Anxiety			PTSD symptoms		
	Event/ total	Weighted prevalence (95% CI)	aOR (95% CI)	Event/ total	Weighted prevalence (95% CI)	aOR (95% CI)	Event/ total	Weighted prevalence (95% CI)	aOR (95% CI)
Initial news outlet									
Traditional media	284/1,149	24.3% (21.8 to 26.9)	1 (ref)	159/1,149	13.4% (11.5 to 15.6)	1 (ref)	141/1,149	11.8% (10.0 to 13.9)	1 (ref)
Online community	66/165	38.4% (30.98 to 46.4)	1.63 (1.12 to 2.38)*	41/165	24.0% (17.9 to 31.4)	1.59 (1.03 to 2.48)*	41/165	23.1% (17.2 to 30.4)	1.55 (0.98 to 2.44)
Social media	95/330	27.8% (23.08 to 33.1)	1.15 (0.84 to 1.58)	44/330	13.3% (9.9 to 17.6)	0.88 (0.58 to 1.34)	57/330	17.4% (13.5 to 22.2)	1.25 (0.83 to 1.90)
Online news	152/532	28.0% (24.18 to 32.3)	1.05 (0.82 to 1.36)	100/532	18.6% (15.3 to 22.4)	1.24 (0.92 to 1.68)	79/532	15.6% (12.5 to 19.4)	1.12 (0.79 to 1.59)
Acquaintance/others	12/44	26.8% (15.2 to 43.0)	0.84 (0.40 to 1.78)	9/44	17.7% (9.1 to 31.6)	1.00 (0.41 to 2.43)	5/44	13.2% (5.1 to 30.2)	0.74 (0.26 to 2.10)
Exposure to graphics									
Not exposed	107/337	31.7% (26.5 to 37.5)	1 (ref)	70/337	20.6% (16.3 to 25.7)	1 (ref)	52/337	17.2% (13.0 to 22.3)	1 (ref)
Exposed	502/1,883	25.8% (23.8 to 27.9)	0.70 (0.53 to 0.94)*	283/1,883	15.4% (13.9 to 17.1)	0.67 (0.48 to 0.93)*	271/1,883	13.8% (12.3 to 15.5)	0.69 (0.47 to 1.01)
Sharing graphics									
Not shared	498/1,924	25.4% (23.4 to 27.5)	1 (ref)	287/1,924	14.7% (13.1 to 16.4)	1 (ref)	240/1,924	12.4% (10.9 to 14.1)	1 (ref)
Shared	111/296	35.2% (29.7 to 41.1)	1.25 (0.93 to 1.68)	66/296	20.3% (16.0 to 25.4)	1.20 (0.86 to 1.69)	83/296	26.7% (21.8 to 32.3)	2.07 (1.46 to 2.93)*

*p<0.05; †age, gender, region of residence, relevance to the victim, initial news outlet, exposure to graphics, sharing graphics, and attribution of responsibility were adjusted. PTSD, posttraumatic stress disorder; CI, confidence interval; OR, odds ratio; aOR, adjusted odds ratio.