



The MAKE Biomarker Discovery for Enhancing antidepressant Treatment Effect and Response (MAKE BETTER) Study: Design and Methodology

Hee-Ju Kang¹, Ju-Wan Kim¹, Seon-Young Kim², Sung-Wan Kim¹,
Hee-Young Shin³, Myung-Geun Shin⁴, and Jae-Min Kim¹ ✉

¹Department of Psychiatry, Chonnam National University Medical School, Gwangju, Republic of Korea

²Mental Health Clinic, Chonnam National University Hwasun Hospital, Hwasun, Republic of Korea

³Department of Biomedical Science, Chonnam National University Medical School, and Clinical Trial Center, Chonnam National University Hospital, Gwangju, Republic of Korea

⁴Department of Laboratory Medicine, Chonnam National University Medical School, Gwangju, Republic of Korea

Psychiatry Investig 2018;15(5):538-545

<https://doi.org/10.30773/pi.2017.10.2>

In the original publication, the schedule of follow-up assessment was incorrect. The description in the study design and recruitment of the Method section has been changed as follows: Eligible patients with depressive disorders who agreed to participate in the MAKE BETTER study were approached for follow-up assessments at 1, 2, 3, 6 and 9 weeks; 3 months; and every 3 months thereafter up to 2 years to determine the short- and long-term treatment outcomes and clinical course of their depressive disorder. Accordingly, the Table 2 has been changed as follows. The authors sincerely regret these missing errors.

Table 2. Schedule of assessment for MAKE BETTER study

	Base-line	1 weeks	2 weeks	3 weeks	6 weeks	9 weeks	3 Month	6 Month	9 Month	12 Month	15 Month	18 Month	21 Month	24 Month	
Clinical assessment															
Socio-demographic characteristics	✓														
Psychological characteristics															
Personality	✓						✓			✓					
Social support	✓						✓			✓					✓
Stress-related status	✓						✓			✓					✓
Outcomes of depression															
Depressive symptom	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Functional disability	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Suicidality	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Quality of life	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Biological assessment															
Blood biomarker	✓									✓					
Salivary cortisol (cortisol study only)	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Electrocardiography	✓									✓					
Resting BP, body mass index	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Treatment related assessment															
Antidepressants, dosage, duration	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Adverse events	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Withdrawal	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓

MAKE BETTER: MAKE Biomarker discovery for Enhancing antidepressants Treatment Effect and Response, BP: blood pressure